

INTERNATIONAL
STANDARD

ISO
3142

Second edition
1997-06-01

**Oil of clove buds [*Syzygium aromaticum*
(L.) Merr. et Perry, syn. *Eugenia*
caryophyllus (Sprengel) Bullock et
S. Harrison]**

Huile essentielle de clous de giroflier [Syzygium aromaticum (L.) Merr. et Perry, syn. Eugenia caryophyllus (Sprengel) Bullock et S. Harrison]

STANDARDSISO.COM : Click to view the full PDF of ISO 3142:1997



Reference number
ISO 3142:1997(E)

Foreword

ISO (the International Organization for Standardization) is a worldwide federation of national standards bodies (ISO member bodies). The work of preparing International Standards is normally carried out through ISO technical committees. Each member body interested in a subject for which a technical committee has been established has the right to be represented on that committee. International organizations, governmental and non-governmental, in liaison with ISO, also take part in the work. ISO collaborates closely with the International Electrotechnical Commission (IEC) on all matters of electrotechnical standardization.

Draft International Standards adopted by the technical committees are circulated to the member bodies for voting. Publication as an International Standard requires approval by at least 75 % of the member bodies casting a vote.

International Standard ISO 3142 was prepared by Technical Committee ISO/TC 54, *Essential oils*.

This second edition cancels and replaces the first edition (ISO 3142:1974), which has been technically revised.

Annexes A et B of this International Standard are for information only.

© ISO 1997

All rights reserved. Unless otherwise specified, no part of this publication may be reproduced or utilized in any form or by any means, electronic or mechanical, including photocopying and microfilm, without permission in writing from the publisher.

International Organization for Standardization
Case Postale 56 • CH-1211 Genève 20 • Switzerland
Internet central@iso.ch
X.400 c=ch; a=400net; p=iso; o=isocs; s=central

Printed in Switzerland

Oil of clove buds [*Syzygium aromaticum* (L.) Merr. et Perry, syn. *Eugenia caryophyllus* (Sprengel) Bullock et S. Harrison]

1 Scope

This International Standard specifies certain characteristics of the oil of clove buds [*Syzygium aromaticum* (L.) Merr. et Perry, syn. *Eugenia caryophyllus* (Sprengel) Bullock and S. Harrison], in order to facilitate assessment of its quality.

2 Normative references

The following standards contain provisions which, through reference in this text, constitute provisions of this International Standard. At the time of publication, the editions indicated were valid. All standards are subject to revision, and parties to agreements based on this International Standard are encouraged to investigate the possibility of applying the most recent editions of the standards indicated below. Members of IEC and ISO maintain registers of currently valid International Standards.

ISO 210:—¹⁾, *Essential oils — General rules for packaging, conditioning and storage.*

ISO 211:—²⁾, *Essential oils — General rules for labelling and marking of containers.*

ISO 212:1973, *Essential oils — Sampling.*

ISO 279:1981, *Essential oils — Determination of relative density at 20 °C (Reference method).*

ISO 280:1976, *Essential oils — Determination of refractive index.*

ISO 292:1981, *Essential oils — Determination of optical rotation.*

ISO 1272:1973, *Essential oils — Determination of phenols content.*

ISO 11024-1:—³⁾, *Essential oils — General guidance on chromatographic profiles — Part 1: Preparation of chromatographic profiles for presentation in standards.*

ISO 11024-2:—³⁾, *Essential oils — General guidance on chromatographic profiles — Part 2: Utilization of chromatographic profiles of a sample of essential oils.*

3 Definition

For the purposes of this International Standard, the following definition applies.

3.1 oil of clove buds: Essential oil obtained by steam distillation of the dried flower buds of clove [*Syzygium aromaticum* (L.) Merr. et Perry, syn. *Eugenia caryophyllus* (Sprengel) Bullock et S. Harrison], of the Myrtaceae family.

1) To be published. (Revision of ISO 210:1961)

2) To be published. (Revision of ISO 211:1961)

3) To be published.

4 Requirements

4.1 Appearance

Clear, mobile liquid, sometimes slightly viscous.

4.2 Colour

From yellow to clear brown.

4.3 Odour

Spicy and characteristic of eugenol.

4.4 Relative density at 20 °C/20 °C

Minimum: 1,042
Maximum: 1,063

4.5 Refractive index at 20 °C

Minimum: 1,528 0
Maximum: 1,538 0

4.6 Optical rotation at 20 °C

Range from –1,5° to 0°.

4.7 Content of total phenolic compounds

Minimum: 85 % (V/V)
Maximum: 93 % (V/V)

4.8 Chromatographic profile

Analysis of the essential oil shall be carried out by gas chromatography. In the chromatogram obtained, the representative and characteristic components shown in table 1 shall be identified. The proportions of these components, indicated by the integrator, shall be as shown in table 1. This constitutes the chromatographic profile of the essential oil.

Table 1 — Chromatographic profile

Component	Minimum %	Maximum %
Eugenol	75	85
β-Caryophyllene	2	7
Eugenyl acetate	8	15

NOTE — The chromatographic profile is normative, contrary to typical chromatograms given for information in annex A.

4.9 Flashpoint

Information on the flashpoint is given in annex B.

5 Sampling

See ISO 212.

Minimum volume of test sample: 25 ml.

NOTE — This volume allows each of the tests specified in this International Standard to be carried out at least once.

6 Test methods

6.1 Relative density at 20 °C/20 °C

See ISO 279.

6.2 Refractive index at 20 °C

See ISO 280.

6.3 Optical rotation at 20 °C

See ISO 592.

6.4 Content of total phenolic compounds

See ISO 1272.

6.5 Chromatographic profile

See ISO 11024-1 and ISO 11024-2.


7 Packaging, labelling, marking and storage

See ISO 210 and ISO 211.

Annex A

(informative)

Typical chromatograms of the essential oil of clove buds (Madagascar origin)




Peak identification

- 1 Eugenol
- 2 β -Caryophyllene
- 3 Eugenyl acetate

Operating conditions

Column: fused silica capillary; length 50 m; diameter 0,25 mm
Stationary phase: polydimethyl siloxane (OV 101)
Oven temperature: from 65 °C to 200 °C, at a rate of 1,5 °C/min
Injector temperature: 250 °C
Detector temperature: 250 °C
Detector: flame ionization
Carrier gas: nitrogen
Volume injected: about 0,2 μ l
Split ratio: 1/100

Figure A.1 — Typical chromatogram taken on an apolar column



Peak identification

- 1 β -Caryophyllene
- 2 Eugenol
- 3 Eugenyl acetate

Operating conditions

Column: fused silica capillary; length 50 m; diameter 0,25 mm
Stationary phase: polyethylene glycol 20 000
Oven temperature: from 65 °C to 200 °C, at a rate of 1,5 °C/min; then at a rate of 1 °C/min up to 230 °C
Injector temperature: 250 °C
Detector temperature: 250 °C
Detector: flame ionization
Carrier gas: nitrogen
Volume injected: about 0,2 μ l
Split ratio: 1/100

Figure A.2 — Typical chromatogram taken on a polar column